

LOWER BOUND ON THE DENSITY OF STATES FOR PERIODIC SCHRÖDINGER OPERATORS

SERGEY MOROZOV, LEONID PARNOVSKI, AND IRINA PCHELINTSEVA

ABSTRACT. We consider Schrödinger operators $-\Delta + V$ in $\mathbb{R}^d$ ($d \geq 2$) with smooth periodic potentials V and prove a uniform lower bound on the density of states for large values of the spectral parameter.

1. INTRODUCTION

Let $H = -\Delta + V$ be a Schrödinger operator in $L_2(\mathbb{R}^d)$ with a smooth periodic potential V . We will assume throughout that $d \geq 2$. The *integrated density of states (IDS)* for H is defined as

$$N(\lambda) := \lim_{L \rightarrow \infty} L^{-d} N(\lambda; H_D^{(L)}), \quad \lambda \in \mathbb{R}. \quad (1.1)$$

Here $H_D^{(L)}$ is the restriction of H to the cube $[0, L]^d$ with the Dirichlet boundary conditions, and $N(\lambda; \cdot)$ is the counting function of the discrete spectrum below λ . For $H_0 := -\Delta$ the IDS can be easily computed explicitly (e.g. using the representation (2.6) below):

$$N_0(\lambda) = (2\pi)^{-d} d^{-1} \omega_d \lambda^{d/2}. \quad (1.2)$$

Here $\omega_d = 2\pi^{d/2}/\Gamma(d/2)$ is the surface area of the unit sphere S^{d-1} in $\mathbb{R}^d$.

The asymptotic behaviour of the function (1.1) for large values of the spectral parameter was recently studied in a number of publications, see [1], [3], and references therein.

Our article concerns the high-energy behaviour of the Radon–Nikodym derivative of IDS

$$g := dN/d\lambda,$$

which is called the *density of states (DOS)* (see [4]). Our main result is that for big values of λ

$$g(\lambda) \geq g_0(\lambda)(1 - o(1)), \quad (1.3)$$

where

$$g_0(\lambda) = dN_0(\lambda)/d\lambda = (2\pi)^{-d} \omega_d \lambda^{(d-2)/2} / 2.$$

We remark that (1.3) should be understood in the sense of measures; in particular, we do not claim that $g(\lambda)$ is everywhere differentiable.

It has been proved in [2] that the spectrum of H contains a semi-axis $[\lambda_0, +\infty)$; this statement is known as the Bethe-Sommerfeld conjecture (see the references in [2] for the history of this problem). This result has an obvious reformulation in terms of IDS: each point $\lambda \geq \lambda_0$ is a point of growth of N . It was also proved in [2] that for each $n \in \mathbb{N}$ and $\varepsilon = \lambda^{-n}$ we have

$$N(\lambda + \varepsilon) - N(\lambda) \ll \varepsilon \lambda^{(d-2)/2}. \quad (1.4)$$

Later, when the second author discussed the results and methods of [2] with Yu. Karpeshina, she suggested that using the technique from that paper, one should be able to prove the opposite bound

$$N(\lambda + \varepsilon) - N(\lambda) \gg \varepsilon \lambda^{(d-2)/2} \quad (1.5)$$

when λ is sufficiently large, not just with $\varepsilon = \lambda^{-n}$ (when the proof is relatively straightforward given [2]), but also uniformly over all $\varepsilon \in (0, 1]$. In this paper we prove that for big λ

$$N(\lambda + \varepsilon) - N(\lambda) \geq \frac{\omega_d}{2(2\pi)^d} \varepsilon \lambda^{(d-2)/2} (1 - o(1)). \quad (1.6)$$

Note that (1.6) implies the claimed bound (1.3).

The proof of (1.6) is heavily based on the technique of [2] and uses various statements proved therein. In order to minimise the size of our paper, we will try to quote as many results as we can from [2], possibly with some minor modifications when necessary.

Acknowledgement. As we have already mentioned, this paper is a result of observations and suggestions made by Yu. Karpeshina; we are very grateful to her for sharing them with us and allowing us to use them. The authors were supported by the EPSRC grant EP/F029721/1.

2. PRELIMINARIES

We study the Schrödinger operator

$$H = -\Delta + V(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^d \quad (2.1)$$

with the potential V being infinitely smooth and periodic with the lattice of periods Λ . We denote the lattice dual to Λ by $\Lambda^\dagger$, fundamental cells of these lattices are denoted by Ω and $\Omega^\dagger$, respectively. We choose $\Omega^\dagger$ to be the first Brillouin zone and introduce

$$Q := \sup \{ |\xi| \mid \xi \in \Omega^\dagger \}. \quad (2.2)$$

Let

$$\mathbf{D} := -i\nabla, \quad \mathbf{D}(\mathbf{k}) := \mathbf{D} + \mathbf{k}. \quad (2.3)$$

The Floquet-Bloch decomposition allows to represent our operator (2.1) as a direct integral (see e.g. [4]):

$$H = \int_{\Omega^\dagger} \oplus H(\mathbf{k}) \, d\mathbf{k}, \quad (2.4)$$

where

$$H(\mathbf{k}) = \mathbf{D}(\mathbf{k})^2 + V(\mathbf{x}) \quad (2.5)$$

is the family of ‘fibre’ operators acting in $L_2(\Omega)$. The domain of each $H(\mathbf{k})$ is the set of periodic functions from $H^2(\Omega)$. The spectrum of H is the union over $\mathbf{k} \in \Omega^\dagger$ of the spectra of the operators (2.5).

We denote by $|\cdot|_\circ$ the surface area Lebesgue measure on the unit sphere S^{d-1} in $\mathbb{R}^d$ and put $\omega_d := |S^{d-1}|_\circ = 2\pi^{d/2}/\Gamma(d/2)$. Finally,

$$N(\lambda) := (2\pi)^{-d} \int_{\Omega^\dagger} \#\{j : \lambda_j(\mathbf{k}) < \lambda\} \, d\mathbf{k} \quad (2.6)$$

is the *integrated density of states* of the operator (2.1). It is known (see e.g. [4]) that the definitions (1.1) and (2.6) are equivalent.

The main result of the paper is

Theorem 2.1. *For sufficiently big λ and any $\varepsilon > 0$ the integrated density of states of H satisfies (1.6).*

By $B(R)$ we denote the ball of radius R centered at the origin. Given two positive functions f and g , we say that $f \gg g$, or $g \ll f$, or $g = O(f)$ if the ratio g/f is bounded. We say $f \asymp g$ if $f \gg g$ and $f \ll g$. Whenever we use O , o , $\gg$, $\ll$, or $\asymp$ notation, the constants involved can depend on d and norms of the potential in various Sobolev spaces H^s ; the same is also the case when we use the expression ‘sufficiently large’. By $\lambda = \rho^2$ we denote a point on the spectral axis. We also denote by v the L_∞ -norm of the potential V , and put $J := [\lambda - 20v, \lambda + 20v]$. Let

$$\mathcal{A} := \left\{ \boldsymbol{\xi} \in \mathbb{R}^d, \quad ||\boldsymbol{\xi}|^2 - \lambda| \leq 40v \right\}. \quad (2.7)$$

Notice that the definition of $\mathcal{A}$ obviously implies that if $\boldsymbol{\xi} \in \mathcal{A}$, then $||\boldsymbol{\xi}| - \rho| \ll \rho^{-1}$.

Any vector $\boldsymbol{\xi} \in \mathbb{R}^d$ can be uniquely decomposed as $\boldsymbol{\xi} = \mathbf{n} + \mathbf{k}$ with $\mathbf{n} \in \Lambda^\dagger$ and $\mathbf{k} \in \Omega^\dagger$. We call $\mathbf{n} = [\boldsymbol{\xi}]$ the ‘integer part’ of $\boldsymbol{\xi}$ and $\mathbf{k} = \{\boldsymbol{\xi}\}$ the ‘fractional part’ of $\boldsymbol{\xi}$.

By $\text{vol}(\cdot)$ we denote the Lebesgue measure in $\mathbb{R}^d$. The identity matrix is denoted by $\mathbf{I}$. For any $h \in L_2(\Omega)$ we introduce its Fourier coefficients

$$h_{\mathbf{n}} := (\text{vol } \Omega)^{-1/2} \int_{\Omega} h(\mathbf{x}) \exp(-i\langle \mathbf{n}, \mathbf{x} \rangle) \, d\mathbf{x}, \quad \mathbf{n} \in \Lambda^\dagger. \quad (2.8)$$

For $\boldsymbol{\xi} \in \mathbb{R}^d \setminus \{\mathbf{0}\}$ we define $r = r(\boldsymbol{\xi}) := |\boldsymbol{\xi}|$ and $\boldsymbol{\xi}' := \boldsymbol{\xi}/|\boldsymbol{\xi}|$. We put

$$R = R(\rho) := \rho^{1/(36d^2(d+2))} \quad (2.9)$$

(so that the condition stated after equation (5.15) in [2] is satisfied). For $j \in \mathbb{N}$ let

$$\Theta'_j := \Lambda^\dagger \cap B(jR) \setminus \{\mathbf{0}\}.$$

Let $M := 5d^2 + 7d$. We introduce the set

$$\mathcal{B} := \left\{ \boldsymbol{\xi} \in \mathcal{A} \mid |\langle \boldsymbol{\xi}, \boldsymbol{\eta}' \rangle| > \rho^{1/2}, \text{ for all } \boldsymbol{\eta} \in \Theta'_{6M} \right\}. \quad (2.10)$$

In other words, $\mathcal{B}$ consists of all points $\boldsymbol{\xi} \in \mathcal{A}$ the projections of which to the directions of all vectors $\boldsymbol{\eta} \in \Theta'_{6M}$ have lengths larger than $\rho^{1/2}$. We also denote $\mathcal{D} := \mathcal{A} \setminus \mathcal{B}$.

In the rest of the section we quote some results from [2] which we will use in this paper. Our approach is slightly different from that of [2]. In particular, we consider arbitrary lattice of periods Λ , not equal to $(2\pi\mathbb{Z})^d$. We also use a different form of the Floquet-Bloch decomposition (so that the operators on fibers (2.5) are defined on the same domain). This leads to several straightforward changes in the formulation of the results from [2]. These changes are:

- (1) The lattices $(2\pi\mathbb{Z})^d$ and $\mathbb{Z}^d$ are replaced by Λ and $\Lambda^\dagger$, respectively. The ‘integer’ and ‘fractional’ parts are now defined with respect to $\Lambda^\dagger$ (see above);
- (2) The matrices $\mathbf{F}$ and $\mathbf{G}$ are replaced by the unit matrix $\mathbf{I}$ throughout;
- (3) The Fourier transform is now defined by (2.8), and the exponentials $e_{\mathbf{m}}$ introduced at the beginning of Section 5 in [2] are redefined as

$$e_{\mathbf{m}}(\mathbf{x}) := (\text{vol } \Omega)^{-1/2} e^{i\langle \mathbf{m}, \mathbf{x} \rangle}, \quad \mathbf{m} \in \Lambda^\dagger;$$

- (4) The operators $H(\mathbf{k})$ are now given by (2.5) on the common domain $\mathfrak{D}$.

The main result we will need follows from Corollary 7.15 of [2]:

Proposition 2.2. *There exist mappings $f, g : \mathcal{A} \rightarrow \mathbb{R}$ which satisfy the following properties:*

- (i) $f(\boldsymbol{\xi})$ is an eigenvalue of $H(\mathbf{k})$ with $\{\boldsymbol{\xi}\} = \mathbf{k}$; $|f(\boldsymbol{\xi}) - |\boldsymbol{\xi}|^2| \leq 2v$. f is an injection (if we count all eigenvalues with multiplicities) and all eigenvalues of $H(\mathbf{k})$ inside J are in the image of f .
- (ii) If $\boldsymbol{\xi} \in \mathcal{A}$, then $|f(\boldsymbol{\xi}) - g(\boldsymbol{\xi})| \leq \rho^{-d-3}$.

(iii) For any $\xi \in \mathcal{B}$

$$g(\xi) = |\xi|^2 + \sum_{j=1}^{2M} \sum_{\eta_1, \dots, \eta_j \in \Theta'_M} \sum_{2 \leq n_1 + \dots + n_j \leq 2M} C_{n_1, \dots, n_j} \langle \xi, \eta_1 \rangle^{-n_1} \dots \langle \xi, \eta_j \rangle^{-n_j}. \quad (2.11)$$

(iv) Let $I = [\mathbf{a}, \mathbf{b}] \subset \mathcal{A}$ be a straight interval of length $L := |\mathbf{b} - \mathbf{a}| \ll \rho^{-1}$. Then there exists an integer vector $\mathbf{n}$ such that $|g(\mathbf{b} + \mathbf{n}) - g(\mathbf{a})| \ll L\rho + \rho^{-d-3}$. Moreover, suppose $\mathbf{m} \neq 0$ is an integer vector such that the interval $I + \mathbf{m}$ is also entirely inside $\mathcal{A}$. Then there exist two different integer vectors $\mathbf{n}_1$ and $\mathbf{n}_2$ such that $|g(\mathbf{b} + \mathbf{n}_1) - g(\mathbf{a})| \ll L\rho + \rho^{-d-3}$ and $|g(\mathbf{b} + \mathbf{n}_2) - g(\mathbf{a} + \mathbf{m})| \ll L\rho + \rho^{-d-3}$.

Remark 2.3. Formula (2.11) implies that

$$\partial g / \partial r(\xi) \asymp \rho, \quad \text{for any } \xi \in \mathcal{B}. \quad (2.12)$$

For each positive $\delta \leq v$ we denote by $\mathcal{A}(\delta)$, $\mathcal{B}(\delta)$, and $\mathcal{D}(\delta)$ the intersections of $g^{-1}([\rho^2 - \delta, \rho^2 + \delta])$ with $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{D}$, respectively.

It is proved in Lemma 8.1 of [2] that

$$\text{vol}(\mathcal{D}(\delta)) \ll \rho^{d-7/3} \delta. \quad (2.13)$$

The following statement (Corollary 8.5 of [2]) gives a sufficient condition for the continuity of f :

Lemma 2.4. *There is a constant C_1 with the following properties. Let*

$$I := \{\xi(t) : t \in [t_{\min}, t_{\max}]\} \subset \mathcal{B}(v).$$

be a straight interval of length $L < \rho^{-1}\delta$. Suppose that there is a point $t_0 \in [t_{\min}, t_{\max}]$ with the property that for each non-zero $\mathbf{n} \in \Lambda^\dagger$ $g(\xi(t_0) + \mathbf{n})$ is either outside the interval

$$\left[g(\xi(t_0)) - C_1 \rho^{-d-3} - C_1 \rho L, g(\xi(t_0)) + C_1 \rho^{-d-3} + C_1 \rho L \right]$$

or not defined. Then $f(\xi(t))$ is a continuous function of t .

By inspection of the proof of Lemma 8.3 of [2] we obtain

Lemma 2.5. *For large enough ρ and $\delta < \rho^{-1}$ the following estimates hold uniformly over $\mathbf{a} \in \Lambda^\dagger \setminus \{\mathbf{0}\}$: if $d \geq 3$,*

$$\text{vol}(\mathcal{B}(\delta) \cap (\mathcal{B}(\delta) + \mathbf{a})) \ll (\delta^2 \rho^{d-3} + \delta \rho^{-d}); \quad (2.14)$$

if $d = 2$,

$$\text{vol} \left(\mathcal{B}(\delta) \cap (\mathcal{B}(\delta) + \mathbf{a}) \right) \begin{cases} \ll \delta^{3/2}, & |\mathbf{a}| \leq 2\rho - 1, \\ \ll \delta^{3/2} + \delta\rho^{-2}, & ||\mathbf{a}| - 2\rho| < 1, \\ = 0, & |\mathbf{a}| \geq 2\rho + 1. \end{cases} \quad (2.15)$$

3. PREVALENCE OF REGULAR DIRECTIONS

Lemma 3.1. *For ρ big enough and*

$$0 < \delta \leq \rho^{-d-3}$$

there exists a set $\mathcal{F} = \mathcal{F}(\rho)$ on the unit sphere S^{d-1} in $\mathbb{R}^d$ with

$$|\mathcal{F}|_o \geq \omega_d(1 - o(1)) \quad (3.1)$$

such that $f(\boldsymbol{\xi})$ is a simple eigenvalue of $H(\{\boldsymbol{\xi}\})$ continuously depending on $r := |\boldsymbol{\xi}|$ for every $\boldsymbol{\xi} = (r, \boldsymbol{\xi}') \in f^{-1}([\rho^2 - \delta, \rho^2 + \delta])$ with $\boldsymbol{\xi}' := \boldsymbol{\xi}/|\boldsymbol{\xi}| \in \mathcal{F}$.

Proof. It is enough to consider $\delta := \rho^{-d-3}$. For each $\boldsymbol{\xi}' \in S^{d-1}$ let

$$I_{\boldsymbol{\xi}'}(\delta) := \{r\boldsymbol{\xi}', r > 0\} \cap \mathcal{B}(\delta). \quad (3.2)$$

Let $\mathcal{F}_1 := \{\boldsymbol{\xi}' \in S^{d-1} \mid I_{\boldsymbol{\xi}'}(\delta) \neq \emptyset, \overline{I_{\boldsymbol{\xi}'}(\delta)} \cap \mathcal{D}(\delta) = \emptyset\}$.

For any $\boldsymbol{\eta} \in \Theta'_{6M}$ the area of the set of points $\boldsymbol{\xi}' \in S^{d-1}$ satisfying

$$|\langle r\boldsymbol{\xi}', \boldsymbol{\eta}' \rangle| \leq \rho^{1/2}$$

is evidently $O(\rho^{-1/2})$ if $r \geq \rho/2$ (the latter is true for all $r\boldsymbol{\xi}' \in \mathcal{A}$). Since the number of elements in Θ'_{6M} is $O(R^d)$, by (2.9) and (2.10) we have

$$|S^{d-1} \setminus \mathcal{F}_1|_o = o(1). \quad (3.3)$$

By definition $\mathcal{B}(\delta) = \mathcal{B} \cap g^{-1}([\rho^2 - \delta, \rho^2 + \delta])$, hence (2.12) implies that for big ρ the length $l_{\boldsymbol{\xi}'}(\delta)$ of $I_{\boldsymbol{\xi}'}(\delta)$ satisfies

$$l_{\boldsymbol{\xi}'}(\delta) \asymp \delta\rho^{-1}, \quad \boldsymbol{\xi}' \in \mathcal{F}_1. \quad (3.4)$$

Let

$$\mathcal{F} := \{\boldsymbol{\xi}' \in \mathcal{F}_1 \mid f \text{ is continuous on } I_{\boldsymbol{\xi}'}(\delta)\},$$

and

$$\mathcal{E}(\delta) := \{\boldsymbol{\xi} \in \mathcal{B}(\delta) \mid \boldsymbol{\xi}' \in \mathcal{F}_1 \setminus \mathcal{F}\}.$$

Lemma 2.4 tells us that for each point $\boldsymbol{\xi} \in \mathcal{E}(\delta)$ there is a non-zero vector $\mathbf{n} \in \Lambda^\dagger$ such that

$$|g(\boldsymbol{\xi} + \mathbf{n}) - g(\boldsymbol{\xi})| \leq C_1(\rho^{-d-3} + \rho l_{\boldsymbol{\xi}'}(\delta)) \ll (\rho^{-d-3} + \delta). \quad (3.5)$$

Since $|g(\boldsymbol{\xi}) - \rho^2| \leq \delta$, this implies

$$|g(\boldsymbol{\xi} + \mathbf{n}) - \rho^2| \leq C_2(\rho^{-d-3} + \delta) =: \delta_1 \ll \rho^{-d-3} = \delta,$$

and thus $\xi + \mathbf{n} \in \mathcal{A}(\delta_1)$; notice that $C_2 > 1$ and so $\delta_1 > \delta$. Therefore, each point $\xi \in \mathcal{E}(\delta)$ also belongs to the set $(\mathcal{A}(\delta_1) - \mathbf{n})$ for a non-zero $\mathbf{n} \in \Lambda^\dagger$; obviously, $|\mathbf{n}| \ll \rho$. In other words,

$$\mathcal{E}(\delta) \subset \bigcup_{\mathbf{n} \in \Lambda^\dagger \cap B(C\rho), \mathbf{n} \neq 0} (\mathcal{A}(\delta_1) - \mathbf{n}) = \bigcup_{\mathbf{n} \neq 0} (\mathcal{B}(\delta_1) - \mathbf{n}) \cup \bigcup_{\mathbf{n} \neq 0} (\mathcal{D}(\delta_1) - \mathbf{n}). \quad (3.6)$$

To proceed further, we need more notation. Denote $\mathcal{D}_0(\delta_1)$ to be the set of all points ν from $\mathcal{D}(\delta_1)$ for which there is no non-zero $\mathbf{n} \in \Lambda^\dagger$ satisfying $\nu - \mathbf{n} \in \mathcal{B}(\delta)$; $\mathcal{D}_1(\delta_1)$ to be the set of all points ν from $\mathcal{D}(\delta_1)$ for which there is a unique non-zero $\mathbf{n} \in \Lambda^\dagger$ satisfying $\nu - \mathbf{n} \in \mathcal{B}(\delta)$; and $\mathcal{D}_2(\delta_1)$ to be the rest of the points from $\mathcal{D}(\delta_1)$ (i.e. $\mathcal{D}_2(\delta_1)$ consists of all points ν from $\mathcal{D}(\delta_1)$ for which there exist at least two different non-zero vectors $\mathbf{n}_1, \mathbf{n}_2 \in \Lambda^\dagger$ satisfying $\nu - \mathbf{n}_j \in \mathcal{B}(\delta)$). Then Lemma 8.7 of [2] implies that we can rewrite (3.6) as

$$\mathcal{E}(\delta) \subset \bigcup_{\mathbf{n} \neq 0} (\mathcal{B}(\delta_1) - \mathbf{n}) \cup \bigcup_{\mathbf{n} \neq 0} (\mathcal{D}_1(\delta_1) - \mathbf{n}). \quad (3.7)$$

This, obviously, implies

$$\mathcal{E}(\delta) \subset \bigcup_{\mathbf{n} \neq 0} \left((\mathcal{B}(\delta_1) - \mathbf{n}) \cap \mathcal{B}(\delta) \right) \cup \bigcup_{\mathbf{n} \neq 0} \left((\mathcal{D}_1(\delta_1) - \mathbf{n}) \cap \mathcal{B}(\delta) \right), \quad (3.8)$$

since $\mathcal{E}(\delta) \subset \mathcal{B}(\delta)$.

The definition of the set $\mathcal{D}_1(\delta_1)$ and (2.13) imply that

$$\begin{aligned} \text{vol} \left(\bigcup_{\mathbf{n} \neq 0} \left((\mathcal{D}_1(\delta_1) - \mathbf{n}) \cap \mathcal{B}(\delta) \right) \right) &\leq \text{vol}(\mathcal{D}_1(\delta_1)) \\ &\leq \text{vol}(\mathcal{D}(\delta_1)) \ll \delta_1 \rho^{d-7/3} \ll \delta \rho^{d-7/3}. \end{aligned} \quad (3.9)$$

For $d \geq 3$ Lemma 2.5, inequality $\delta < \delta_1$, and the fact that the union in (3.8) consists of no more than $C\rho^d$ terms imply

$$\text{vol} \left(\bigcup_{\mathbf{n} \neq 0} \left((\mathcal{B}(\delta_1) - \mathbf{n}) \cap \mathcal{B}(\delta) \right) \right) \ll \rho^d (\delta_1^2 \rho^{d-3} + \delta_1 \rho^{-d}) \ll \delta (\rho^{d-6} + 1). \quad (3.10)$$

For $d = 2$ we obtain by Lemma 2.5

$$\begin{aligned}
& \text{vol} \left(\bigcup_{\mathbf{n} \in \Lambda^\dagger \setminus \{\mathbf{0}\}} \left(\mathcal{B}(\delta) \cap (\mathcal{B}(\delta_1) + \mathbf{n}) \right) \right) \\
& \leq \sum_{\substack{\mathbf{n} \in \Lambda^\dagger \setminus \{\mathbf{0}\} \\ |\mathbf{n}| \leq 2\rho - 1}} \text{vol} \left(\mathcal{B}(\delta) \cap (\mathcal{B}(\delta_1) + \mathbf{n}) \right) \\
& + \sum_{\substack{\mathbf{n} \in \Lambda^\dagger \setminus \{\mathbf{0}\} \\ ||\mathbf{n}| - 2\rho| < 1}} \text{vol} \left(\mathcal{B}(\delta) \cap (\mathcal{B}(\delta_1) + \mathbf{n}) \right) \\
& \ll \delta_1^{3/2} \rho^2 + \rho(\delta_1^{3/2} + \delta_1 \rho^{-2}) \ll \delta \rho^{-1/2},
\end{aligned} \tag{3.11}$$

where we have used that

$$\#\left\{ \mathbf{n} \in \Lambda^\dagger \mid ||\mathbf{n}| - 2\rho| < 1 \right\} \ll \rho.$$

Applying (3.9), (3.10), and (3.11) to (3.8) we obtain for all $d \geq 2$

$$\text{vol } \mathcal{E}(\delta) \ll \delta \rho^{d-7/3}. \tag{3.12}$$

By definition,

$$\mathcal{E}(\delta) = \bigcup_{\boldsymbol{\xi}' \in \mathcal{F}_1 \setminus \mathcal{F}} I_{\boldsymbol{\xi}'}(\delta).$$

Hence by (3.4)

$$|\mathcal{F}_1 \setminus \mathcal{F}|_o \ll \delta^{-1} \rho^{2-d} \text{vol } \mathcal{E}(\delta). \tag{3.13}$$

Combining (3.12) and (3.13) we conclude that for big ρ

$$|\mathcal{F}_1 \setminus \mathcal{F}|_o = o(1). \tag{3.14}$$

We have

$$|S^{d-1} \setminus \mathcal{F}|_o = |S^{d-1} \setminus \mathcal{F}_1|_o + |\mathcal{F}_1 \setminus \mathcal{F}|_o. \tag{3.15}$$

Substituting (3.3) and (3.14) into (3.15) we obtain (3.1).

Now we notice that for every $\boldsymbol{\xi}' \in \mathcal{F}$ the interval $I_{\boldsymbol{\xi}'}(\delta)$ has the following property: for each point $\boldsymbol{\xi} \in I_{\boldsymbol{\xi}'}(\delta)$ and each non-zero vector $\mathbf{n} \in \Lambda^\dagger$ such that $\boldsymbol{\xi} + \mathbf{n} \in \mathcal{A}$ we have $|g(\boldsymbol{\xi} + \mathbf{n}) - g(\boldsymbol{\xi})| > 2\rho^{-d-3}$. This implies $f(\boldsymbol{\xi} + \mathbf{n}) - f(\boldsymbol{\xi}) \neq 0$. Therefore, $f(\boldsymbol{\xi})$ is a simple eigenvalue of $H(\{\boldsymbol{\xi}\})$ for each $\boldsymbol{\xi} \in I_{\boldsymbol{\xi}'}(\delta)$. The lemma is proved. $\square$

4. SOME PROPERTIES OF OPERATORS ON THE FIBERS

For $m \in \mathbb{R}$ let

$$V^{(m)} := \left(\sum_{\mathbf{n} \in \Lambda^\dagger} |\mathbf{n}|^{2m} |V_{\mathbf{n}}|^2 \right)^{1/2}.$$

Since V is smooth, $V^{(m)}$ is finite for any $m \geq 0$. Recall that Q is defined by (2.2).

Lemma 4.1. *Fix $m \in \mathbb{N}$ and $\varkappa \in (0, 1)$. For $\mathbf{k} \in \Omega^\dagger$ let ψ be a normalized eigenfunction of $H(\mathbf{k})$:*

$$H(\mathbf{k})\psi = \zeta\psi \quad (4.1)$$

with the eigenvalue

$$\zeta \geq \max \{36Q^2\varkappa^{-2}, (1 + m\varkappa)^{2/(d-1)}\varkappa^{-2d/(d-1)}\}. \quad (4.2)$$

Then there exists $M_m = M_m(d, \Lambda, V) \in \mathbb{R}_+$ such that for all $\mathbf{n} \in \Lambda^\dagger$ with

$$|\mathbf{n}| \geq (1 + m\varkappa)\sqrt{\zeta} \quad (4.3)$$

the Fourier coefficients of ψ satisfy

$$|\psi_{\mathbf{n}}| < M_m\varkappa^{-m}|\mathbf{n}|^{-(3m+1)/2}. \quad (4.4)$$

Proof. We proceed by induction. Suppose that either $m = 1$, or $m > 1$ and the statement is proved for $m - 1$. Substituting the Fourier series

$$\psi(\mathbf{x}) = (\text{vol } \Omega)^{-1/2} \sum_{\mathbf{n} \in \Lambda^\dagger} \psi_{\mathbf{n}} \exp(i\langle \mathbf{n}, \mathbf{x} \rangle), \quad \mathbf{x} \in \Omega$$

into (4.1) and equating the coefficients at $\exp(i\langle \mathbf{n}, \mathbf{x} \rangle)$ on both sides, we obtain by (2.5):

$$|\mathbf{n} + \mathbf{k}|^2 \psi_{\mathbf{n}} + \sum_{\mathbf{l} \in \Lambda^\dagger} V_{\mathbf{n}-\mathbf{l}} \psi_{\mathbf{l}} = \zeta \psi_{\mathbf{n}}. \quad (4.5)$$

Since $|\mathbf{k}| \leq Q$, by (4.2) and (4.3) we have

$$2|\mathbf{n}||\mathbf{k}| \leq \varkappa|\mathbf{n}|^2/6 + 6\varkappa^{-1}Q^2 \leq \varkappa|\mathbf{n}|^2/3. \quad (4.6)$$

For $\varkappa \in (0, 1)$, it follows from (4.3) that

$$|\mathbf{n}|^2 - \zeta \geq (1 - (1 + \varkappa)^{-2})|\mathbf{n}|^2 = \varkappa(2 + \varkappa)(1 + \varkappa)^{-2}|\mathbf{n}|^2 \geq \varkappa|\mathbf{n}|^2/2. \quad (4.7)$$

Combining (4.6) and (4.7) we obtain

$$|\mathbf{n} + \mathbf{k}|^2 - \zeta \geq |\mathbf{n}|^2 - 2|\mathbf{n}||\mathbf{k}| - \zeta \geq \varkappa|\mathbf{n}|^2/6,$$

and thus by (4.5)

$$|\psi_{\mathbf{n}}| < 6\varkappa^{-1}|\mathbf{n}|^{-2} \sum_{\mathbf{l} \in \Lambda^\dagger} |V_{\mathbf{n}-\mathbf{l}} \psi_{\mathbf{l}}|. \quad (4.8)$$

If $m = 1$ we estimate the sum on the r. h. s. by $V^{(0)}$ using Cauchy–Schwarz inequality (since ψ is normalized) and obtain (4.4) with $M_1 := 6V^{(0)}$.

If $m > 1$, we estimate

$$\sum_{\mathbf{l} \in \Lambda^\dagger: |\mathbf{l} - \mathbf{n}| \leq |\mathbf{n}|^{1/d}} |V_{\mathbf{n}-\mathbf{l}} \psi_1| \leq \sup_{\mathbf{m}: |\mathbf{m}| \geq |\mathbf{n}| - |\mathbf{n}|^{1/d}} |\psi_{\mathbf{m}}| \sum_{\mathbf{l} \in \Lambda^\dagger: |\mathbf{l}| \leq |\mathbf{n}|^{1/d}} |V_{\mathbf{l}}|. \quad (4.9)$$

By (4.3), (4.2), and monotonicity of the function $q(t) = t - t^{1/d}$ for $t > 1$ we have

$$|\mathbf{n}| - |\mathbf{n}|^{1/d} \geq (1 + m\kappa)\sqrt{\zeta} - ((1 + m\kappa)\sqrt{\zeta})^{1/d} \geq (1 + (m-1)\kappa)\sqrt{\zeta}.$$

We can thus apply the induction hypothesis obtaining

$$\sup_{\mathbf{m}: |\mathbf{m}| \geq |\mathbf{n}| - |\mathbf{n}|^{1/d}} |\psi_{\mathbf{m}}| \leq \kappa^{1-m} M_{m-1} (1 - |\mathbf{n}|^{(1-d)/d})^{1-3m/2} |\mathbf{n}|^{1-3m/2}. \quad (4.10)$$

Since $\kappa \in (0, 1)$, by (4.3) and (4.2) we have

$$|\mathbf{n}| \geq (1 + m\kappa)\sqrt{\zeta} \geq (\kappa^{-1} + m)^{d/(d-1)} > 2^{d/(d-1)},$$

thus

$$(1 - |\mathbf{n}|^{(1-d)/d})^{1-3m/2} < 2^{3m/2-1}. \quad (4.11)$$

Let

$$W := \sup_{r>1} r^{-d} \# \{ \mathbf{l} \in \Lambda^\dagger : |\mathbf{l}| \leq r \}.$$

Clearly, $W < \infty$. Then by Cauchy–Schwarz inequality

$$\sum_{\mathbf{l} \in \Lambda^\dagger: |\mathbf{l}| \leq |\mathbf{n}|^{1/d}} |V_{\mathbf{l}}| \leq W^{1/2} V^{(0)} |\mathbf{n}|^{1/2}. \quad (4.12)$$

Substituting (4.10), (4.11), and (4.12) into (4.9) we get

$$\sum_{\mathbf{l} \in \Lambda^\dagger: |\mathbf{l} - \mathbf{n}| \leq |\mathbf{n}|^{1/d}} |V_{\mathbf{n}-\mathbf{l}} \psi_1| < 2^{3m/2-1} \kappa^{1-m} W^{1/2} V^{(0)} M_{m-1} |\mathbf{n}|^{3(1-m)/2}. \quad (4.13)$$

On the other hand, since $\|\psi\| = 1$, by Cauchy–Schwarz we have

$$\begin{aligned} \sum_{\mathbf{l} \in \Lambda^\dagger: |\mathbf{l} - \mathbf{n}| > |\mathbf{n}|^{1/d}} |V_{\mathbf{n}-\mathbf{l}} \psi_1| &< |\mathbf{n}|^{3(1-m)/2} \sum_{\mathbf{l} \in \Lambda^\dagger: |\mathbf{l}| > |\mathbf{n}|^{1/d}} |\mathbf{l}|^{3(m-1)d/2} |V_{\mathbf{l}}| |\psi_{\mathbf{n}-\mathbf{l}}| \\ &\leq |\mathbf{n}|^{3(1-m)/2} \left(\sum_{\mathbf{l} \in \Lambda^\dagger: |\mathbf{l}| > |\mathbf{n}|^{1/d}} |\mathbf{l}|^{3(m-1)d} |V_{\mathbf{l}}|^2 \right)^{1/2} \leq V^{(3(m-1)d/2)} |\mathbf{n}|^{3(1-m)/2}. \end{aligned} \quad (4.14)$$

Inserting (4.13) and (4.14) into (4.8) we obtain (4.4) with

$$M_m := 6(2^{3m/2-1} W^{1/2} V^{(0)} M_{m-1} + V^{(3(m-1)d/2)}).$$

□

Lemma 4.2. *For any $\eta \in (0, 1)$ there exists $\zeta_0 > 0$ such that if $\zeta(\mathbf{k}) \geq \zeta_0$ is a simple eigenvalue of $H(\mathbf{k})$ for some $\mathbf{k} \in \Omega^\dagger$ then*

$$|\nabla_{\mathbf{k}}\zeta| \leq 2(1 + \eta)\sqrt{\zeta}. \quad (4.15)$$

Proof. Let $\psi(\mathbf{k})$ be the eigenfunction corresponding to $\zeta(\mathbf{k})$ with

$$\|\psi(\mathbf{k})\| = 1. \quad (4.16)$$

Then

$$\nabla_{\mathbf{k}}\zeta(\mathbf{k}) = \nabla_{\mathbf{k}}(\psi(\mathbf{k}), H(\mathbf{k})\psi(\mathbf{k})) = \left(\psi(\mathbf{k}), (\nabla_{\mathbf{k}}H(\mathbf{k}))\psi(\mathbf{k}) \right). \quad (4.17)$$

By (2.5) and (2.3),

$$\nabla_{\mathbf{k}}H(\mathbf{k}) = 2\mathbf{D}(\mathbf{k}).$$

Substituting this into (4.17) we obtain:

$$|\nabla_{\mathbf{k}}\zeta(\mathbf{k})| \leq 2\|\mathbf{D}(\mathbf{k})\psi(\mathbf{k})\| = 2\left(\sum_{\mathbf{n} \in \Lambda^\dagger} |\mathbf{n} + \mathbf{k}|^2 |\psi_{\mathbf{n}}(\mathbf{k})|^2 \right)^{1/2}. \quad (4.18)$$

Let

$$m := [(d+1)/3] + 1 \quad (4.19)$$

and

$$\varkappa := \eta/(2m+1). \quad (4.20)$$

We put

$$\zeta_0 := \max \{ 36Q^2\varkappa^{-2}, (1 + m\varkappa)^{2/(d-1)}\varkappa^{-2d/(d-1)} \} \quad (4.21)$$

and assume that $\zeta := \zeta(\mathbf{k}) \geq \zeta_0$. Since by (2.2) $|\mathbf{k}| \leq Q$, by (4.16), (4.21), and (4.20) we have

$$\sum_{|\mathbf{n}| < (1+m\varkappa)\sqrt{\zeta}} |\mathbf{n} + \mathbf{k}|^2 |\psi_{\mathbf{n}}(\mathbf{k})|^2 < (1 + (m+1/6)\varkappa)^2 \zeta < (1 + \eta/2)^2 \zeta. \quad (4.22)$$

For $|\mathbf{n}| \geq (1 + m\varkappa)\sqrt{\zeta}$ we apply Lemma 4.1 obtaining

$$\sum_{|\mathbf{n}| \geq (1+m\varkappa)\sqrt{\zeta}} |\mathbf{n} + \mathbf{k}|^2 |\psi_{\mathbf{n}}(\mathbf{k})|^2 \leq M_m^2 \varkappa^{-2m} \sum_{|\mathbf{n}| \geq (1+m\varkappa)\sqrt{\zeta}} |\mathbf{n} + \mathbf{k}|^2 |\mathbf{n}|^{-3m-1}. \quad (4.23)$$

By (4.19) the r. h. s. of (4.23) is finite and is $O(\zeta^{-1/2})$. Thus, increasing ζ_0 if necessary, by (4.18), (4.22), and (4.23) we obtain (4.15). $\square$

5. PROOF OF THEOREM 2.1

It is enough to prove

Theorem 5.1. *For any $\alpha \in (0, 1)$ there exists $\rho_0 > 0$ big enough such that for all $\rho \geq \rho_0$*

$$N(\rho^2 + \delta) - N(\rho^2 - \delta) \geq (1 - \alpha)(2\pi)^{-d} \omega_d \delta \rho^{d-2} \quad (5.1)$$

for any

$$0 < \delta \leq \rho^{-d-3}. \quad (5.2)$$

Indeed, the original statement of Theorem 2.1 can be obtained by partitioning of the interval $[\lambda, \lambda + \varepsilon]$ into subintervals with lengths not exceeding $2\lambda^{(-d-3)/2}$ and adding up estimates (5.1) on this subintervals (with ρ^2 being respective middle points).

Proof. We first express the growth of IDS in terms of the function f of Proposition 2.2:

$$N(\rho^2 + \delta) - N(\rho^2 - \delta) = (2\pi)^{-d} \text{vol} (f^{-1}[\rho^2 - \delta, \rho^2 + \delta]). \quad (5.3)$$

We can write

$$\text{vol} (f^{-1}[\rho^2 - \delta, \rho^2 + \delta]) = \int_{S^{d-1}} \int_0^\infty \chi(r, \xi') r^{d-1} dr d\xi', \quad (5.4)$$

where χ is the indicator function of $f^{-1}([\rho^2 - \delta, \rho^2 + \delta])$. To obtain a lower bound we can restrict the integration in (5.4) to $\xi' \in \mathcal{F}$ defined in Lemma 3.1. Then for any $\eta \in (0, 1)$ there exists $\rho_0 > 0$ such that for any $\rho \geq \rho_0$ we have

$$|\mathcal{F}|_\circ \geq (1 - \eta)\omega_d, \quad (5.5)$$

and for any $\xi' \in \mathcal{F}$ the support of $\chi(\cdot, \xi')$ contains an interval $[r_1, r_2]$ with

$$(1 - \eta)\rho \leq r_1 < r_1 + (1 - \eta)\rho^{-1}\delta \leq r_2. \quad (5.6)$$

Indeed, the first inequality in (5.6) follows from Proposition 2.2(ii),(iii). The last inequality in (5.6) follows from Lemmata 3.1 and 4.2.

Thus for all $\rho \geq \rho_0$ by (5.5) and (5.6) we obtain

$$\begin{aligned} \int_{S^{d-1}} \int_0^\infty \chi(r, \xi') r^{d-1} dr d\xi' &\geq \int_{\mathcal{F}} (1 - \eta)^d \rho^{d-2} \delta d\xi' \\ &\geq (1 - \eta)^{d+1} \omega_d \rho^{d-2} \delta, \end{aligned} \quad (5.7)$$

Combining (5.3), (5.4), and (5.7), and choosing η small enough we arrive at (5.1). The theorem is proved. $\square$

REFERENCES

1. Yulia E. Karpeshina, *On the density of states for the periodic Schrödinger operator*, Ark. Mat. **38** (2000), no. 1, 111–137. MR MR1749362 (2001g:47088)
2. Leonid Parnowski, *Bethe-Sommerfeld conjecture*, Ann. Henri Poincaré **9** (2008), no. 3, 457–508. MR MR2419769
3. Leonid Parnowski and Roman Shterenberg, *Asymptotic expansion of the integrated density of states of a two-dimensional periodic Schrödinger operator*, Invent. Math. **176** (2009), no. 2, 275–323.
4. Michael Reed and Barry Simon, *Methods of modern mathematical physics. IV. Analysis of operators*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1978. MR MR0493421 (58 #12429c)

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON, WC1E 6BT, UK